

Vector Bundles on Products of Varieties with n -blocks Collections

Edoardo Ballico and Francesco Malaspina

Università di Trento
38050 Povo (TN), Italy
e-mail: ballico@science.unitn.it

Politecnico di Torino
Corso Duca degli Abruzzi 24, 10129 Torino, Italy
e-mail: malaspina@calvino.polito.it

April 1, 2008

1

Abstract

Here we consider the product of varieties with n -blocks collections . We give some cohomological splitting conditions for rank 2 bundles. A cohomological characterization for vector bundles is also provided. The tools are Beilinson's type spectral sequences generalized by Costa and Miró-Roig. Moreover we introduce a notion of Castelnuovo-Mumford regularity on a product of finitely many projective spaces and smooth quadric hypersurfaces in order to prove two splitting criteria for vector bundle with arbitrary rank.

Introduction

A well known result by Horrocks (see [9]) characterizes the vector bundles without intermediate cohomology on a projective space as direct sum of line bundles. This criterion fails on more general varieties. In fact there exist non-split vector bundles on X without intermediate cohomology. These bundles are called ACM bundles.

On a quadric hypersurface $\mathcal{Q}_n$ there is a theorem that classifies all the ACM bundles (see [12]) as direct sums of line bundles and spinor bundles (up to a twist - for generalities about spinor bundles see [16]).

Ottaviani has generalized Horrocks criterion to quadrics and Grassmannians giving cohomological splitting conditions for vector bundles (see [15] and [17]).

The starting point of this note is [6] where Laura Costa and Rosa Maria Miró-Roig give a new proof of Horrocks and Ottaviani's criteria by using different techniques. Beilinson's Theorem was stated in 1978 and since then it has become a major tool in classifying vector bundles over projective spaces. Beilinson's spectral sequence was generalized by Kapranov (see [10])

¹Mathematics Subject Classification 2000: 14F05, 14J60.

keywords: n -blocks Collections; vector bundles; Castelnuovo-Mumford regularity.

and [11]) to hyperquadrics and Grassmannians and by Costa and Miró-Roig (see [6]) to any smooth projective variety of dimension n with a n -block collection.

Our aim is to use these results and techniques in order to give some cohomological splitting conditions for rank 2 bundles on products of varieties with n -blocks collections.

We also give some Cohomological characterization for vector bundles. Horrocks in [9] gives a cohomological characterization of p -differentials over $\mathbb{P}^n$.

Ancona and Ottaviani in [1] give this characterization on quadrics and Costa and Miró-Roig on multiprojective spaces. Here we give some different characterization on quadrics and we extend the result by Costa and Miró-Roig on any product of varieties with n -blocks collections.

In the last section we specialize on a product X of finitely many projective spaces and smooth quadric hypersurfaces. In [2] and [3] we introduced a notion of Castelnuovo-Mumford regularity on quadric hypersurfaces and multiprojective spaces. We will give a suitable definition of regularity on such a product X in order to prove splitting criteria for vector bundle with arbitrary rank. Let E be a vector bundle on X . We will give two criteria which says when E is (up to a twist) a direct sum of $\mathcal{O}$ or the tensor product of pull-backs of spinor bundles on the quadric factors of X (see Theorems 4.11 and 4.12).

We want to thank Claudio Fontanari for the helpful discussions and Laura Costa and Rosa Maria Miró-Roig for having showed us theirs preprints, which were the starting point of our work.

1 Preliminaries

Throughout the paper X will be a smooth projective variety defined over the complex numbers $\mathbb{C}$ and we denote by $\mathcal{D} = D^b(\mathcal{O}_X - \text{mod})$ the derived category of bounded complexes of coherent sheaves of $\mathcal{O}_X$ -modules.

For the notations we refer to [6].

Now we give the definition of n -block collection in order to introduce a Beilinson's type spectral sequence generalized:

Definition 1.1. *An exceptional collection $(F_0, F_1, \dots, F_m)$ of objects of $\mathcal{D}$ (see [6] Definition 2.1.) is a block if $\text{Ext}_{\mathcal{D}}^i(F_j, F_k) = 0$ for any i and $j \neq k$.*

An n -block collection of type $(\alpha_0, \alpha_1, \dots, \alpha_n)$ of objects of $\mathcal{D}$ is an exceptional collection

$$(\mathcal{E}_0, \mathcal{E}_1, \dots, \mathcal{E}_m) = (E_1^0, \dots, E_{\alpha_0}^0, E_1^1, \dots, E_{\alpha_1}^1, \dots, E_1^n, \dots, E_{\alpha_n}^n)$$

such that all the subcollections $\mathcal{E}_i = (E_1^i, \dots, E_{\alpha_i}^i)$ are blocks.

Notation 1.2. *If $\mathcal{F}$ is a bundle on X and*

$$(\mathcal{E}_0, \mathcal{E}_1, \dots, \mathcal{E}_m) = (E_1^0, \dots, E_{\alpha_0}^0, E_1^1, \dots, E_{\alpha_1}^1, \dots, E_1^n, \dots, E_{\alpha_n}^n)$$

is an n -block collection, we denote by $\overline{\mathcal{E}}_j$ the direct sum of all the bundle in $\mathcal{E}_j$,

$$\overline{\mathcal{E}}_j = \bigoplus_{i=1}^{\alpha_j} E_i^j.$$

So we have

$$\mathcal{F} \otimes \overline{\mathcal{E}}_j = \bigoplus_{i=1}^{\alpha_j} (\mathcal{F} \otimes E_i^j).$$

Example 1.3. $(\mathcal{O}_{\mathbb{P}^n}(-n), \mathcal{O}_{\mathbb{P}^n}(-n+1), \dots, \mathcal{O}_{\mathbb{P}^n})$ is an n -block collection of type $(1, 1, \dots, 1)$ on $\mathbb{P}^n$ (see [6] Example 2.3.(1)).

Example 1.4. Let us consider a smooth quadric hypersurface $\mathcal{Q}_n$ in $\mathbb{P}^{n+1}$.

We use the unified notation Σ_* meaning that for even n both the spinor bundles Σ_1 and Σ_2 are considered, and for n odd, the spinor bundle Σ . We follow the notation of [6] so the spinor bundles are twisted by 1 with respect to those of [16] ($\Sigma_* = \mathcal{S}_*(1)$) $(\mathcal{E}_0, \mathcal{O}(-n+1), \dots, \mathcal{O}(-1), \mathcal{O})$, where $\mathcal{E}_0 = (\Sigma_*(-n))$, is an n -block collection of type $(1, 1, \dots, 1)$ if n is odd, and of type $(2, 1, \dots, 1)$ if n is even (see [6] Example 3.4.(2)). Moreover we can have several n -block collections:

$$\sigma_j = (\mathcal{O}(j), \dots, \mathcal{O}(n-1), \mathcal{E}_{n-j}, \mathcal{O}(n+1), \dots, \mathcal{O}(n+j-1))$$

where $\mathcal{E}_{n-j} = (\Sigma_*(n-1))$ and $1 \leq j \leq n$ (see [7] Proposition 4.4).

Example 1.5. Let $X = G(k, n)$ be the Grassmannian of k -dimensional subspaces of the n -dimensional vector space. Assume $k > 1$. We have the canonical exact sequence

$$0 \rightarrow \mathcal{S} \rightarrow \mathcal{O}^n \rightarrow \mathcal{Q} \rightarrow 0,$$

where $\mathcal{S}$ denote the tautological rank k bundle and $\mathcal{Q}$ the quotient bundle.

Denote by $A(k, n)$ the set of locally free sheaves $\Sigma^\alpha \mathcal{S}$ where α runs over diagrams fitting inside a $k \times (n-k)$ rectangle. We have $\Sigma^{(p, 0, \dots, 0)} \mathcal{S} = \mathcal{S}^p \mathcal{S}$, $\Sigma^{(1, 1, 0, \dots, 0)} \mathcal{S} = \wedge^2 \mathcal{S} = \mathcal{S}(1)$ and $\Sigma^{(p_1+t, p_2+t, \dots, p_k+t)} \mathcal{S} = \Sigma^{(p_1, p_2, \dots, p_k)} \mathcal{S}(t)$.

$A(k, n)$ can be totally ordered in such a way that we obtain a $k(n-k)$ -block collection $\sigma = (\mathcal{E}_0, \dots, \mathcal{E}_{k(n-k)})$ by packing in the same block $\mathcal{E}_r$ the bundles $\Sigma^\alpha \mathcal{S}$ with $|\alpha| = k(n-k) - r$ (see [6] Example 3.4.(1)).

Example 1.6. Let $X = \mathbf{P}^{n_1} \times \dots \times \mathbf{P}^{n_r}$ be a multiprojective space and $d = n_1 + \dots + n_r$. For any $0 \leq j \leq r$, denote by $\mathcal{E}_j$ the collection of all line bundles on X

$$\mathcal{O}_X(a_1^j, a_2^j, \dots, a_r^j)$$

with $-n_i \leq a_i^j \leq 0$ and $\sum_{i=1}^r a_i^j = j - d$. Using the Künneth formula we prove that $\mathcal{E}_j$ is a block and that

$$(\mathcal{E}_0, \dots, \mathcal{E}_d)$$

is a d -block collection of line bundles on X (see [6] Example 3.4.(3)).

Beilinson's Theorem was generalized by Costa and Miró-Roig to any smooth projective variety of dimension n with an n -block collection of coherent sheaves which generates $\mathcal{D}$ (see [6] Theorem 3.6.).

2 Products of n -blocks Collections

Let X, Y be two smooth projective varieties of dimension n and m . Let $(\mathcal{G}_0, \dots, \mathcal{G}_n)$, $\mathcal{G}_i = (G_0^i, \dots, G_{\alpha_i}^i)$ be a n -block collection for X and $(\mathcal{F}_0, \dots, \mathcal{F}_m)$, $\mathcal{F}_i = (F_0^i, \dots, F_{\beta_i}^i)$ a m -block collection for Y .

Notation 2.1. We denote by $\mathcal{G}_i \boxtimes \mathcal{F}_j$ the set of all the bundles $G_k \boxtimes F_m$ on $X \times Y$ such that $G_k \in \mathcal{G}_i$ and $F_m \in \mathcal{F}_j$.

For any $0 \leq k \leq n + m$, we define $\mathcal{E}_k = \mathcal{G}_i \boxtimes \mathcal{F}_j$ where $i + j = k$.

Lemma 2.2. $(\mathcal{E}_0, \dots, \mathcal{E}_{n+m})$ is a $(m + n)$ -block collection for $X \times Y$ and it generates the derived category $\mathcal{D}$.

Proof. By [5] Proposition 3.4. we have that $(\mathcal{E}_0, \dots, \mathcal{E}_{n+m})$ is a strongly exceptional collection and by [5] Proposition 4.16. it generates the derived category $\mathcal{D}$.

We only need to prove that each $\mathcal{E}_i$ is a block. Let $G_j \boxtimes F_k$ and $G_l \boxtimes F_m$ two different bundles of $\mathcal{E}_i$ we have to show that

$$\text{Ext}^0(G_j \boxtimes F_k, G_l \boxtimes F_m) = \text{Ext}^0(G_j \boxtimes G_l) \otimes \text{Ext}^0(F_k \boxtimes F_m) = 0.$$

Now, since $j + k = l + m = i$ and $(j, k) \neq (l, m)$, we have $j < l$ that means $\text{Ext}^0(G_j \boxtimes G_l) = 0$ or $k < m$ that means $\text{Ext}^0(F_k \boxtimes F_m) = 0$.

All the other vanishing comes from the fact that $(\mathcal{E}_0, \dots, \mathcal{E}_{n+m})$ is a strongly exceptional collection. $\square$

Remark 2.3. Let $X = X_1 \times \dots \times X_r$ where $\dim X_i = n_i$ and $d = n_1 + \dots + n_r$. For every X_i let us consider the n_i -blocks collection $(\mathcal{G}_0^i, \dots, \mathcal{G}_{n_i}^i)$. Let assume that for any $i = 1, \dots, r$, $\mathcal{G}_{n_i}^i = \{\mathcal{O}\}$.

For any $0 \leq k \leq r$, denote by $\mathcal{E}_k$ the collection of all bundles on X

$$\mathcal{G}_{j_1}^1 \boxtimes \dots \boxtimes \mathcal{G}_{j_r}^r$$

with $\sum_{i=1}^r j_i = k$.

$(\mathcal{E}_0, \dots, \mathcal{E}_d)$ is a d -block collection on X and it generates the derived category $\mathcal{D}$.

Example 2.4. Let $X = \mathcal{Q}_{n_1} \times \dots \times \mathcal{Q}_{n_r}$ be a multiquadric space and $d = n_1 + \dots + n_r$. Let us assume $n_i \geq 3$ for every i .

For every $\mathcal{Q}_{n_i}$ let us consider the n_i -block collection $(\mathcal{G}_0^i, \dots, \mathcal{G}_{n_i}^i)$, where $\mathcal{G}_0^i = (\Sigma_*(-n_i))$ and $\mathcal{G}_j^i = \mathcal{O}(-n_i + j)$ for $1 \leq j \leq n_i$.

Now we define a d -block collection on X .

For any $0 \leq k \leq r$, denote by $\mathcal{E}_k$ the collection of all bundles on X

$$\mathcal{G}_{j_1}^1 \boxtimes \dots \boxtimes \mathcal{G}_{j_r}^r$$

with $\sum_{i=1}^r j_i = k$.

On $X = X_1 \times \dots \times X_r$ we can use the Beilinson's type spectral sequences generalized by Costa and Miró-Roig in order to prove splitting criteria.

We want to apply [6] Proposition 4.1. for a bundle E on X . We need for any i , $0 \leq i \leq d - 1$,

$$H^i(X, E \otimes \mathcal{E}_{d-i-1}) = 0.$$

In particular

$$H^0(X, E \otimes \mathcal{E}_{d-1}) = 0.$$

Here $\mathcal{E}_d = (\mathcal{O}_X(0, \dots, 0))$ but $\mathcal{E}_{d-1}$ is more complicated so we need the following definition:

Definition 2.5. A bundle E on $X = X_1 \times \cdots \times X_r$ is normalized if $H^0(X, E) \neq 0$ and $H^0(X, E \otimes \mathcal{E}_{d-1}) = 0$.

In particular on multiquadrics E is normalized if $H^0(X, E) \neq 0$ and $H^0(X, E \otimes \mathcal{O}_X(a_1, \dots, a_r)) = 0$ when $a_1, \dots, a_r$ are non-positive integers not all vanishing.

Now we can prove the following result (cfr [13]):

Proposition 2.6. Let $X = X_1 \times \cdots \times X_r$ as in Remark 2.3. Let E be a normalized bundle on X . If, for any i , $1 \leq i \leq d-1$,

$$H^i(X, E \otimes \mathcal{G}_{j_1}^1 \boxtimes \cdots \boxtimes \mathcal{G}_{j_r}^r) = 0$$

when $\sum_{h=1}^r j_h = i+1$, then E contains $\mathcal{O}_X$ as a direct summand.

Proof. Since E is normalized $H^0(X, E) \neq 0$ but $H^0(X, E \otimes \mathcal{E}_{d-1}) = 0$. So we can apply [6] Proposition 4.1. for E and conclude that E contains $\overline{\mathcal{E}_d} \cong \mathcal{O}$ as a direct summand. $\square$

Remark 2.7. The above Proposition cannot become a splitting criterion because it is not possible to iterate the above argument. In fact we cannot obtain normalized bundles by tensoring only with $\mathcal{O}(t, \dots, t)$. We have a splitting criteria only for rank two bundles

Corollary 2.8. Let $X = X_1 \times \cdots \times X_r$ as in Remark 2.3. Let E be a rank two normalized bundle on X . If, for any i , $1 \leq i \leq d-1$,

$$H^i(X, E \otimes \mathcal{G}_{j_1}^1 \boxtimes \cdots \boxtimes \mathcal{G}_{j_r}^r) = 0$$

when $\sum_{h=1}^r j_h = i+1$, then E splits as a direct sum of line bundles.

Remark 2.9. In the last section we will see that, when $X = \mathbf{P}^{n_1} \times \cdots \times \mathbf{P}^{n_s} \times \mathcal{Q}_{m_1} \times \cdots \times \mathcal{Q}_{m_q}$ by using different techniques, it is possible to prove splitting criteria for bundles with arbitrary rank.

3 Cohomological Characterization of Vector Bundles

Horrocks in [9] gives a cohomological characterization of p -differentials over $\mathbb{P}^n$.

Ancona and Ottaviani in [1] extend this characterization on quadrics:

they show that, if $\mathcal{F}$ is a coherent sheaf on $\mathcal{Q}_n$ and for some $t \in \mathbb{Z}$ and some $0 < j < n$

$$H^i(\mathcal{F}(t-i-1)) = 0 \text{ for } j \leq i \leq n-2$$

$$H^i(\mathcal{F}(t-i+1)) = 0 \text{ for } 1 \leq i \leq j$$

$$H^{n-1}(\mathcal{F} \otimes \Sigma(t-n)) = 0 \text{ for any spinor bundle } \Sigma,$$

then $\mathcal{F}$ contains $\psi_j^{h^i(\mathcal{F}(t-j))}$ as direct summand (see [1] Theorem 6.3.).

Moreover, let $\mathcal{F}$ be a coherent sheaf on $\mathcal{Q}_n$ and suppose that for some $t \in \mathbb{Z}$

$$H^i(\mathcal{F}(t-i-1)) = 0 \text{ for } 0 \leq i \leq n-2, \text{ and}$$

$$H^{n-1}(\mathcal{F} \otimes \Sigma(t-n)) = H^{n-1}(\mathcal{F}(t-n)) = 0, \text{ then}$$

for n odd $\mathcal{F}$ contains $\Sigma^\vee(-t+1)^{h^{n-1}(\mathcal{F} \otimes \Sigma(t-n))}$ as direct summand

for n even $\mathcal{F}$ contains $\Sigma_1^\vee(-t+1)^{h^{n-1}(\mathcal{F} \otimes \Sigma_2(t-n))} \oplus \Sigma_2^\vee(-t+1)^{h^{n-1}(\mathcal{F} \otimes \Sigma_1(t-n))}$ as direct summand

(see [1] Theorem 6.7.).

Remark 3.1. By using different n -block collections on $\mathcal{Q}_n$ we can obtain different cohomological characterizations.

Let us consider for instance the n -block collection:

$$(\mathcal{O}(-n+1), \mathcal{E}_1, \mathcal{O}(-n+2), \dots, \mathcal{O})$$

where $\mathcal{E}_1 = (\Sigma_*(-n+1))$ and its right dual

$$(\mathcal{O}, \psi_1^\vee, \dots, \psi_{n-2}^\vee, \mathcal{H}_{n-1}, \mathcal{O}(1))$$

where $\mathcal{H}_{n-1} = (\Sigma_*)$, (see [7] Proposition 4.4).

Let $\mathcal{F}$ be a coherent sheaf on $\mathcal{Q}_n$ and suppose that for some $t \in \mathbb{Z}$

$H^i(\mathcal{F}(t-i+1)) = 0$ for $1 \leq i \leq n-1$, and

$H^{n-1}(\mathcal{F}(t-n+1)) = 0$, then

for n odd $\mathcal{F}$ contains $\Sigma^\vee(-t)^{h^{n-1}(\mathcal{F} \otimes \Sigma(t-n+1))}$ as direct summand

for n even $\mathcal{F}$ contains $\Sigma_1^\vee(-t)^{h^{n-1}(\mathcal{F} \otimes \Sigma_2(t-n+1))} \oplus \Sigma_2^\vee(-t)^{h^{n-1}(\mathcal{F} \otimes \Sigma_1(t-n+1))}$ as direct summand.

Moreover if $\mathcal{F}$ is a coherent sheaf on $\mathcal{Q}_n$ and for some $t \in \mathbb{Z}$ and some $0 < j < n-1$

$H^i(\mathcal{F}(t-i-1)) = 0$ for $j \leq i \leq n-2$

$H^i(\mathcal{F}(t-i+1)) = 0$ for $1 \leq i \leq j$

$H^{n-2}(\mathcal{F} \otimes \Sigma(t-n+1)) = 0$ for any spinor bundle Σ ,

$H^{n-1}(\mathcal{F}(t-n+1)) = 0$

then $\mathcal{F}$ contains $\psi_j^{h^i(\mathcal{F}(t-j))}$ as direct summand.

Proof. Our cohomological conditions are corresponding to those of [6] Proposition 4.8 so we can conclude that E contains as a direct summand the term E_1^{-jj} of the spectral sequence defined in [6] Theorem 3.6.

E_1^{-jj} is $\Sigma^\vee(-t)^{h^{n-1}(\mathcal{F} \otimes \Sigma(t-n+1))}$ or $\Sigma_1^\vee(-t)^{h^{n-1}(\mathcal{F} \otimes \Sigma_2(t-n+1))} \oplus \Sigma_2^\vee(-t)^{h^{n-1}(\mathcal{F} \otimes \Sigma_1(t-n+1))}$ if $j = n-1$ and it is $\psi_j^{h^i(\mathcal{F}(t-j))}$ if $0 < j < n-1$. $\square$

Laura Costa and Rosa Maria Miró-Roig give a cohomological characterization on multi-projective spaces ([6] Proposition 4.10 and Theorem 4.11) and grassmannians (see [6] Corollary 4.12.).

Let us consider now in general the products of varieties with an n -block collection. We have the following result:

Lemma 3.2. Let X, Y be two smooth projective varieties of dimension n and m . Let $(\mathcal{G}_0, \dots, \mathcal{G}_n)$ and $(\mathcal{F}_0, \dots, \mathcal{F}_m)$ be an n -block collection for X and an m -block collection for Y . Let $({}'\mathcal{G}_0, \dots, {}'\mathcal{G}_n)$ and $({}'\mathcal{F}_0, \dots, {}'\mathcal{F}_m)$ be their right dual collections (see [6] Proposition 3.9.).

For any $0 \leq k \leq n+m$, we define $'\mathcal{E}_k = {}'\mathcal{G}_i \otimes {}'\mathcal{F}_j$ where $i+j=k$.

Then $({}'\mathcal{E}_0, \dots, {}'\mathcal{E}_{n+m})$ is the right dual $(m+n)$ -block collection of $(\mathcal{E}_0, \dots, \mathcal{E}_{n+m})$ on $X \times Y$.

Proof. $({}'\mathcal{E}_0, \dots, {}'\mathcal{E}_{n+m})$ is an $(m+n)$ -block collection by the above lemma.

In order to prove that it is the right dual collection of $(\mathcal{E}_0, \dots, \mathcal{E}_{n+m})$ we only need to show that it verifies the orthogonality conditions (see [6] (3.7) and (3.8)).

For every couple of bundles $\mathcal{G} \boxtimes \mathcal{F}$ and $'\mathcal{G} \boxtimes {}'\mathcal{F}$ on $X \times Y$,

$$H^\alpha(X \times Y, \mathcal{G} \boxtimes \mathcal{F} \otimes {}'\mathcal{G} \boxtimes {}'\mathcal{F}) = \bigoplus_{\alpha_1 + \alpha_2 = \alpha} H^{\alpha_1}(X, \mathcal{G} \otimes {}'\mathcal{G}) \otimes H^{\alpha_2}(Y, \mathcal{F} \otimes {}'\mathcal{F}).$$

So the orthogonality conditions of $(\mathcal{G}_0, \dots, \mathcal{G}_n)$ and $(\mathcal{F}_0, \dots, \mathcal{F}_m)$ on X and Y imply those of $(\mathcal{E}_0, \dots, \mathcal{E}_{n+m})$ on $X \times Y$. $\square$

Remark 3.3. Let $X = X_1 \times \dots \times X_r$ where $\dim X_i = n_i$ and $d = n_1 + \dots + n_r$. For every X_i let us consider the n_i -blocks collection $(\mathcal{G}_0^i, \dots, \mathcal{G}_{n_i}^i)$ and the right dual n_i -blocks collection $(\mathcal{G}_0^i, \dots, \mathcal{G}_{n_i}^i)$.

For any $0 \leq k \leq r$, denote by $\mathcal{E}_k$ the collection of all bundles on X

$$\mathcal{G}_{j_1}^1 \boxtimes \dots \boxtimes \mathcal{G}_{j_r}^r$$

with $\sum_{i=1}^r j_i = k$.

$(\mathcal{E}_0, \dots, \mathcal{E}_d)$ is the right dual d -block collection of $(\mathcal{E}_0, \dots, \mathcal{E}_d)$.

The block $\mathcal{E}_{d-j}$ is the orthogonal block of the block $\mathcal{E}_j$ because it contains all the orthogonal bundles of the bundles in $\mathcal{E}_j$.

Example 3.4. Let us consider $X = \mathbf{P}^{n_1} \times \dots \times \mathbf{P}^{n_r} \times \mathcal{Q}_{m_1} \times \dots \times \mathcal{Q}_{m_s}$ with $d = n_1 + \dots + n_r + m_1 + \dots + m_s$. On every $\mathbf{P}^{n_1}$ we consider the n_i -block collection:

$$\mathcal{G}^{n_i} = (\mathcal{O}(-n_i), \mathcal{O}(-n_i + 1), \dots, \mathcal{O})$$

and its right dual

$$(\mathcal{O}, \mathbb{T}(-1), \dots, \wedge^{n_i} \mathbb{T}(-n_i))$$

and on every $\mathcal{Q}_{m_i}$ we consider the m_i -block collection:

$$\mathcal{H}^{m_i} = (\mathcal{O}(-m_i + 1), \mathcal{H}_1^{m_i}, \mathcal{O}(-m_i + 2), \dots, \mathcal{O})$$

where $\mathcal{H}_1^{m_i} = (\Sigma_*(-m_i + 1))$ and its right dual

$$(\mathcal{O}, \psi_1^\vee, \dots, \psi_{m_i-2}^\vee, \mathcal{H}_{m_i-1}^{m_i}, \mathcal{O}(1))$$

where $\mathcal{H}_{m_i-1}^{m_i} = (\Sigma_*)$.

We can obtain a d -block collection

$$(\mathcal{E}_0, \dots, \mathcal{E}_d)$$

as in Remark 2.3.

For any $\mathcal{O}(t_1) \boxtimes \dots \boxtimes \mathcal{O}(t_r) \boxtimes \mathcal{O}(k_1) \boxtimes \dots \boxtimes \mathcal{O}(k_h) \boxtimes \Sigma_*(-m_{h+1} - 1) \boxtimes \dots \boxtimes \Sigma_*(-m_s - 1) \in \mathcal{E}_{d-k}$ and any $0 \leq k \leq d$ ($0 \leq h \leq s$).

$$R_{\mathcal{E}_d \dots \mathcal{E}_{d-k+1}} \mathcal{O}(t_1) \boxtimes \dots \boxtimes \mathcal{O}(t_r) \boxtimes \mathcal{O}(k_1) \boxtimes \dots \boxtimes \mathcal{O}(k_h) \boxtimes \Sigma_*(-m_{h+1} - 1) \boxtimes \dots \boxtimes \Sigma_*(-m_s - 1) = \\ \wedge^{-t_1} \mathbb{T}(t_1) \boxtimes \dots \boxtimes \wedge^{-t_r} \mathbb{T}(t_r) \boxtimes \psi_{-k_1}^\vee(k_1) \boxtimes \dots \boxtimes \psi_{-k_h}^\vee(k_h) \boxtimes \Sigma_* \boxtimes \dots \boxtimes \Sigma_*$$

Proposition 3.5. Let $X = X_1 \times \dots \times X_r$ where $\dim X_i = n_i$ and $d = n_1 + \dots + n_r$. Let $(\mathcal{E}_0, \dots, \mathcal{E}_d)$ be the d -block collection with right dual $(\mathcal{E}_0, \dots, \mathcal{E}_d)$ as in Remark 3.3. We denote by r_j the rank of $\overline{\mathcal{E}_j}$.

Let E be a bundle on X .

Assume there exists j , $0 < j < n$ such that for any $-n \leq p \leq -j - 1$

$$H^{-p-1}(X, E \otimes \overline{\mathcal{E}_{p+n}}) = 0$$

and for any $j + 1 \leq p \leq 0$

$$H^{-p+1}(X, E \otimes \overline{\mathcal{E}_{p+n}}) = 0.$$

If rank $E = r_{d-j}$, then

$$E \cong \overline{\mathcal{E}_{d-j}}.$$

Proof. Our cohomological conditions are corresponding to those of [6] Proposition 4.8 so we can conclude that E contains as a direct summand the term E_1^{-jj} of the spectral sequence defined in [6] Theorem 3.6.

Now, since $\text{rank } E = r_{d-j}$, then

$$E \cong \overline{\mathcal{E}_{d-j}}.$$

□

Example 3.6. Let us consider $X = \mathbf{P}^3 \times \mathcal{Q}_3$.

On $\mathbf{P}^3$ we have the collection $(\mathcal{O}(-3), \dots, \mathcal{O})$ and the right dual $(\mathcal{O}, \mathbb{T}(-1), \dots, \wedge^3 \mathbb{T}(-3))$ and on $\mathcal{Q}_3$ we have the collection $(\Sigma(-3), \mathcal{O}(-2), \dots, \mathcal{O})$ and the right dual $(\mathcal{O}, \psi_1^\vee, \psi_2^\vee, \Sigma)$. We can obtain a 6-block collection

$$(\mathcal{E}_0, \dots, \mathcal{E}_6)$$

as in Remark 2.3.

Let E be a rank 12 bundle such that

$$H^1(X, E \otimes \overline{\mathcal{E}_6}) = H^1(X, E) = 0$$

and for any $1 \leq i \leq 5$

$$H^i(X, E \otimes \overline{\mathcal{E}_{5-i}}) = 0.$$

Then

$$E \cong (\mathbb{T}(1) \boxtimes \mathcal{O}) \oplus (\mathcal{O} \boxtimes \psi_1) \cong \mathbb{T}(1) \boxtimes \psi_1.$$

4 Regularity on $\mathbf{P}^{n_1} \times \dots \times \mathbf{P}^{n_s} \times \mathcal{Q}_{m_1} \times \dots \times \mathcal{Q}_{m_q}$

In [2] we introduced the following definition of regularity on $\mathcal{Q}_n$ (cfr [2] Definition 2.1 and Proposition 2.4):

Definition 4.1. A coherent sheaf F on $\mathcal{Q}_n$ ($n \geq 2$) is said to be m - Q regular if $H^i(F(m-i)) = 0$ for $i = 1, \dots, n-1$, $H^{n-1}(F(m) \otimes \Sigma_*(-n+1)) = 0$ and $H^n(F(m-n+1)) = 0$.

We will say Q regular in order to 0- Q regular.

In [3] we introduced the following definition of regularity on $\mathbf{P}^{n_1} \times \dots \times \mathbf{P}^{n_s}$ (cfr [3] Definition 4.1):

Definition 4.2. A coherent sheaf F on $\mathbf{P}^{n_1} \times \dots \times \mathbf{P}^{n_s}$ is said to be $(p_1, \dots, p_s)$ -regular if, for all $i > 0$,

$$H^i(F(p_1, \dots, p_s) \otimes \mathcal{O}(k_1, \dots, k_s)) = 0$$

whenever $k_1 + \dots + k_s = -i$ and $-n_j \leq k_j \leq 0$ for any $j = 1, \dots, s$.

Now we want to introduce a notion of regularity on $\mathbf{P}^{n_1} \times \dots \times \mathbf{P}^{n_s} \times \mathcal{Q}_{m_1} \times \dots \times \mathcal{Q}_{m_q}$. Let us consider $X = \mathbf{P}^n \times \mathcal{Q}_m$.

Definition 4.3. On $\mathbf{P}^n$ we consider the n -block collection:

$$(\mathcal{E}_0, \dots, \mathcal{E}_n) = (\mathcal{O}(-n), \mathcal{O}(-n+1), \dots, \mathcal{O})$$

and on $\mathcal{Q}_m$ we consider the m -block collection:

$$(\mathcal{G}_0, \dots, \mathcal{G}_m) = (\mathcal{O}(-m+1), \mathcal{G}_1, \dots, \mathcal{O})$$

where $\mathcal{G}_1 = (\Sigma_*(-m+1))$.

A coherent sheaf F on X is said to be (p, p') -regular if, for all $i > 0$,

$$H^i(F(p, p') \otimes \mathcal{E}_{n-j} \boxtimes \mathcal{G}_{m-k}) = 0$$

whenever $j + k = i$, $-n \leq -j \leq 0$ and $-m \leq -k \leq 0$.

Remark 4.4. If $m = 2$ Definition 4.3 coincides with Definition 4.2 on $\mathbf{P}^n \times \mathbf{P}^1 \times \mathbf{P}^1$. In fact the 2-block collection on $\mathcal{Q}_2$ is

$$(\mathcal{O}(-1), \{\Sigma_1(-1), \Sigma_2(-1)\}, \mathcal{O}) = (\mathcal{O}(-1, -1), \{\mathcal{O}(-1, 0), \mathcal{O}(0, -1)\}, \mathcal{O}).$$

Remark 4.5. If $m = 0$ we can identify X with $\mathbf{P}^n$ and the sheaf $F(k, k')$ with $F(k)$. Under this identification F is (p, p') -regular in the sense of Definition 4.3, if and only if F is p -regular in the sense of Castelnuovo-Mumford.

In fact, let $i > 0$, $H^i(F(p, p') \otimes \mathcal{E}_{n-j} \boxtimes \mathcal{G}_{m-k}) = H^i(F(p-j)) = 0$ whenever $j + k = i$, $-n \leq -j \leq 0$ and $-m \leq -k \leq 0$ if and only if $H^i(F(p-j)) = 0$ whenever $-i \leq -j \leq 0$ if and only if $H^i(F(p'-i)) = 0$.

If $n = 0$ we can identify X with $\mathcal{Q}_m$ and the sheaf $F(k, k')$ with $F(k')$. Under this identification F is (p, p') -regular in the sense of Definition 4.3, if and only if F is p' -Qregular on $\mathcal{Q}_m$. In fact, let $i > 0$, $H^i(F(p, p') \otimes \mathcal{E}_{n-j} \boxtimes \mathcal{G}_{m-k}) = H^i(F(p') \otimes \mathcal{G}_{m-k}) = 0$ whenever $j + k = i$, $-n \leq -j \leq 0$ and $-m \leq -k \leq 0$ if and only if $H^i(F(p') \otimes \mathcal{G}_{m-i}) = 0$ if and only if F is p' -Qregular.

Lemma 4.6. (1) Let H be a generic hyperplane of $\mathbf{P}^n$. If F is a regular coherent sheaf on $X = \mathbf{P}^n \times \mathcal{Q}_m$, then $F|_{L_1}$ is regular on $L_1 = H \times \mathcal{Q}_m$.

(2) Let H' be a generic hyperplane of $\mathcal{Q}_m$. If F is a regular coherent sheaf on $X = \mathbf{P}^n \times \mathcal{Q}_m$, then $F|_{L_2}$ is regular on $L_2 = \mathbf{P}^n \times H'$.

Proof. (1) We follow the proof of [8] Lemma 2.6.. We get this exact cohomology sequence:

$$H^i(F(-j, 0) \otimes \mathcal{O} \boxtimes \mathcal{G}_{m-k}) \rightarrow H^i(F|_{L_1}(-j, 0) \otimes \mathcal{O} \boxtimes \mathcal{G}_{m-k}) \rightarrow H^i(F(-j-1, 0) \otimes \mathcal{O} \boxtimes \mathcal{G}_{m-k})$$

If $j + k = i$, $-n \leq -j \leq 0$ and $-m \leq -k \leq 0$, we have also $-n-1 \leq j-1 \leq 0$, so the first and the third groups vanish by hypothesis. Then also the middle group vanishes and $F|_{L_1}$ is regular.

(2) We have to deal also with the spinor bundles. First assume m even, say $m = 2l$. We have $\Sigma_1|_{\mathcal{Q}_{m-1}} \cong \Sigma_2|_{\mathcal{Q}_{m-1}} \cong \Sigma$. Let us consider the exact sequences

$$0 \rightarrow \mathcal{O}(-j) \boxtimes \Sigma_1(-m) \rightarrow \mathcal{O}(-j) \boxtimes \mathcal{O}(-m+1)^{2^l} \rightarrow \mathcal{O}(-j) \boxtimes \Sigma_2(-m+1) \rightarrow 0$$

tensored by F .

Let $i \geq m-1$ and $j = m-1-j$. Since $H^i(F \otimes \mathcal{O}(-j) \boxtimes \Sigma_2(-m+1)) = H^i(F \otimes \mathcal{E}_{n-j} \boxtimes \mathcal{G}_1) = 0$ and $H^{i+1}(F(-j, -m+1)) = H^{i+1}(F \otimes \mathcal{E}_{n-j} \boxtimes \mathcal{G}_0) = 0$, we also have

$$H^{i+1}(F \otimes \mathcal{O}(-j) \boxtimes \Sigma_1(-m)) = 0.$$

From the exact sequences

$$0 \rightarrow \mathcal{O}(-j) \boxtimes \Sigma_1(-m+1) \rightarrow \mathcal{O}(-j) \boxtimes \Sigma_1(-m+2) \rightarrow \mathcal{O}(-j) \boxtimes \Sigma_1|_{\mathcal{Q}_{m-1}}(-m+2) \rightarrow 0$$

tensored by F , we get

$$H^i(F(-j, 0) \boxtimes \Sigma_1(-m+1)) \rightarrow H^i(F(-j, 0) \boxtimes \Sigma_1|_{\mathcal{Q}_{m-1}}(-m+1)) \rightarrow H^{i+1}(F(-j, 0) \boxtimes \Sigma_1(-m))$$

If $i \geq m-1$ and $j = m-1-j$, the first and the third groups vanish by hypothesis. Then also the middle group vanishes.

Assume now m odd, say $m = 2l+1$. We have $\Sigma_1|_{\mathcal{Q}_{m-1}} \cong \Sigma_1 \oplus \Sigma_2$. We can consider the exact sequences

$$0 \rightarrow \mathcal{O}(-j) \boxtimes \Sigma(-m) \rightarrow \mathcal{O}(-j) \boxtimes \mathcal{O}(-m+1)^{2^{l+1}} \rightarrow \mathcal{O}(-j) \boxtimes \Sigma(-m+1) \rightarrow 0$$

tensored by F . Then we argue as above.

All the others vanishing in Definition 4.3 can be proved as in (1) and we can conclude that $F|_{L_2}$ is regular. □

Proposition 4.7. *Let F be a regular coherent sheaf on $X = \mathbf{P}^n \times \mathcal{Q}_m$ then*

1. $F(p, p')$ is regular for $p, p' \geq 0$.

2. $H^0(F(k, k'))$ is spanned by

$$H^0(F(k-1, k')) \otimes H^0(\mathcal{O}(1, 0))$$

if $k-1, k' \geq 0$; and it is spanned by

$$H^0(F(k, k'-1)) \otimes H^0(\mathcal{O}(0, 1))$$

if $k, k'-1 \geq 0$ and $m > 2$.

Proof. (1) We will prove part (1) by induction. Let F be a regular coherent sheaf, we want show that also $F(1, 0)$ is regular. We follow the proof of [8] Proposition 2.7.

Consider the exact cohomology sequence:

$$H^i(F(-j, 0) \otimes \mathcal{O} \boxtimes \mathcal{G}_{m-k}) \rightarrow H^i(F(-j+1, 0) \otimes \mathcal{O} \boxtimes \mathcal{G}_{m-k}) \rightarrow H^i(F|_{L_1}(-j+1, 0) \otimes \mathcal{O} \boxtimes \mathcal{G}_{m-k})$$

If $j+k=i$, $-n \leq -j \leq 0$ and $-m \leq -k \leq 0$, so the first and the third groups vanish by hypothesis. Then also the middle group vanishes.

A symmetric argument shows the vanishing for $F(0, 1)$. We only have to check the vanishing involving the spinor bundles.

$$H^i(F(-j, 0) \boxtimes \Sigma_*(-m+1)) \rightarrow H^i(F(-j, 1) \boxtimes \Sigma_*(-m+1)) \rightarrow H^i(F(-j, 1) \boxtimes \Sigma_*|_{\mathcal{Q}_{m-1}}(-m+1))$$

If $i \geq m-1$ and $j = m-1-j$, the first and the third groups vanish by hypothesis. Then also the middle group vanishes.

(2) We will follow the proof of [8] Proposition 2.8.

We consider the following diagram:

$$\begin{array}{ccc} H^0(F(k-1, k')) \otimes H^0(\mathcal{O}(1, 0)) & \xrightarrow{\sigma} & H^0(F|_{L_1}(k-1, k')) \otimes H^0(\mathcal{O}_{L_1}(1, 0)) \\ \downarrow \mu & & \downarrow \tau \\ H^0(F(k, k')) & \xrightarrow{\nu} & H^0(F|_{L_1}(k, k')) \end{array}$$

Note that σ is surjective if $k-1, k' \geq 0$ because $H^1(F(k-2, k')) = 0$ by regularity.

Moreover also τ is surjective by (2) for $F|_{L_1}$.

Since both σ and τ are surjective, we can see as in [14] page 100 that μ is also surjective.

In order to prove that $H^0(F(k, k'))$ is spanned by $H^0(F(k, k'-1)) \otimes H^0(\mathcal{O}(0, 1))$ if $k, k'-1 \geq 0$, we can use a symmetric argument since for $m > 2$ the spinor bundles are not involved in the proof. $\square$

Remark 4.8. *If F is a regular coherent sheaf on $X = \mathbf{P}^n \times \mathcal{Q}_m$ ($m > 2$) then it is globally generated.*

In fact by the above proposition we have the following surjections:

$$H^0(F) \otimes H^0(\mathcal{O}(1, 0)) \otimes H^0(\mathcal{O}(0, 1)) \rightarrow H^0(F(1, 0)) \otimes H^0(\mathcal{O}(0, 1)) \rightarrow H^0(F(1, 1)),$$

and so the map

$$H^0(F) \otimes H^0(\mathcal{O}(1, 1)) \rightarrow H^0(F(1, 1))$$

is a surjection.

Moreover we can consider a sufficiently large twist l such that $F(l, l)$ is globally generated.

The commutativity of the diagram

$$\begin{array}{ccc} H^0(F) \otimes H^0(\mathcal{O}(l, l)) \otimes \mathcal{O} & \rightarrow & H^0(F(l, l)) \otimes \mathcal{O} \\ \downarrow & & \downarrow \\ H^0(F) \otimes \mathcal{O}(l, l) & \rightarrow & F(l, l) \end{array}$$

yields the surjectivity of $H^0(F) \otimes \mathcal{O}(l, l) \rightarrow F(l, l)$, which implies that F is generated by its sections.

If $m = 2$, then F is spanned by Remark 4.4 and [3] Remark 2.6.

Now we generalize Definition 4.3:

Definition 4.9. *Let us consider $X = \mathbf{P}^{n_1} \times \cdots \times \mathbf{P}^{n_s} \times \mathcal{Q}_{m_1} \times \cdots \times \mathcal{Q}_{m_q}$.*

On $\mathbf{P}^{n_j}$ (where $j = 1, \dots, s$) we consider the n_j -block collections:

$$(\mathcal{E}_0^j, \dots, \mathcal{E}_n^j) = (\mathcal{O}(-n_j), \mathcal{O}(-n_j + 1), \dots, \mathcal{O})$$

and on $\mathcal{Q}_{m_l}$ (where $l = 1, \dots, q$) we consider the m_l -block collections:

$$(\mathcal{G}_0^l, \dots, \mathcal{G}_m^l) = (\mathcal{O}(-m_l + 1), \mathcal{G}_1^l, \dots, \mathcal{O})$$

where $\mathcal{G}_1^l = (\Sigma_*(-m_l + 1))$.

A coherent sheaf F on X is said to be $(p_1, \dots, p_{s+q})$ -regular if, for all $i > 0$,

$$H^i(F(p_1, \dots, p_{s+q}) \otimes \mathcal{E}_{n_1-k_1}^1 \boxtimes \cdots \boxtimes \mathcal{E}_{n_s-k_s}^s \boxtimes \mathcal{G}_{m_1-h_1}^1 \boxtimes \cdots \boxtimes \mathcal{G}_{m_q-h_q}^q) = 0$$

whenever $k_1 + \cdots + k_s + h_1 + \cdots + h_q = i$, $-n_j \leq -k_j \leq 0$ for any $j = 1, \dots, s$ and $-m_l \leq -h_l \leq 0$ for any $l = 1, \dots, q$.

Remark 4.10. All the above properties (in particular Remark 4.8) can be proved (by using exactly the same arguments) also for this extension of Definition 4.3 (cfr [3])

We use our notion of regularity in order to proving some splitting criterion on $X = \mathbf{P}^{n_1} \times \cdots \times \mathbf{P}^{n_s} \times \mathcal{Q}_{m_1} \times \cdots \times \mathcal{Q}_{m_q}$.

Theorem 4.11. Let E be a rank r vector bundle on $X = \mathbf{P}^{n_1} \times \cdots \times \mathbf{P}^{n_s} \times \mathcal{Q}_{m_1} \times \cdots \times \mathcal{Q}_{m_q}$ ($m_1, \dots, m_q > 2$). Set $d = n_1 + \cdots + n_s + m_1 + \cdots + m_q$. Then the following conditions are equivalent:

1. for any $i = 1, \dots, d-1$ and for any integer t ,

$$H^i(E(t, \dots, t) \otimes \mathcal{E}_{n_1-k_1}^1 \boxtimes \cdots \boxtimes \mathcal{E}_{n_s-k_s}^s \boxtimes \mathcal{G}_{m_1-h_1}^1 \boxtimes \cdots \boxtimes \mathcal{G}_{m_q-h_q}^q) = 0$$

whenever $k_1 + \dots + k_s + h_1 + \cdots + h_q = i$, $-n_j \leq k_j \leq 0$ for any $j = 1, \dots, s$ and $-m_l \leq -h_l \leq 0$ for any $l = 1, \dots, q$.

2. There are r integer $t_1, \dots, t_r$ such that $E \cong \bigoplus_{i=1}^r \mathcal{O}(t_i, \dots, t_i)$.

Proof. (1) $\Rightarrow$ (2). Let assume that t is an integer such that $E(t, \dots, t)$ is regular but $E(t-1, \dots, t-1)$ not.

By the definition of regularity and (1) we can say that $E(t-1, \dots, t-1)$ is not regular if and only if

$$H^d(E(t-1, \dots, t-1) \otimes \mathcal{O}(-n_1, \dots, -n_s, -m_1+1, \dots, -m_q+1)) \neq 0.$$

By Serre duality we have that $H^0(E^\vee(-t, \dots, -t)) \neq 0$.

Now since $E(t, \dots, t)$ is globally generated by Remark 4.8 and $H^0(E^\vee(-t, \dots, -t)) \neq 0$ we can conclude that $\mathcal{O}$ is a direct summand of $E(t, \dots, t)$.

By iterating these arguments we get (2).

(2) $\Rightarrow$ (1). By Künneth formula for any $i = 1, \dots, m+n-1$ and for any integer t ,

$$H^i(\mathcal{O}(t, \dots, t) \otimes \mathcal{E}_{n_1-k_1}^1 \boxtimes \cdots \boxtimes \mathcal{E}_{n_s-k_s}^s \boxtimes \mathcal{G}_{m_1-h_1}^1 \boxtimes \cdots \boxtimes \mathcal{G}_{m_q-h_q}^q) = 0$$

whenever $k_1 + \dots + k_s + h_1 + \cdots + h_q = i$, $-n_j \leq k_j \leq 0$ for any $j = 1, \dots, s$ and $-m_l \leq -h_l \leq 0$ for any $l = 1, \dots, q$.

Then $\mathcal{O}$ satisfies all the conditions in (1). □

Theorem 4.12. Let E be a rank r vector bundle on $X = \mathbf{P}^{n_1} \times \cdots \times \mathbf{P}^{n_s} \times \mathcal{Q}_{m_1} \times \cdots \times \mathcal{Q}_{m_q}$ ($m_1, \dots, m_q > 2$). Set $d = n_1 + \cdots + n_s + m_1 + \cdots + m_q$. Then the following conditions are equivalent:

1. for any $i = 1, \dots, d-1$ and for any integer t ,

$$H^i(E(t, \dots, t) \otimes \mathcal{E}_{n_1-k_1}^1 \boxtimes \cdots \boxtimes \mathcal{E}_{n_s-k_s}^s \boxtimes \mathcal{G}_{m_1-h_1}^1 \boxtimes \cdots \boxtimes \mathcal{G}_{m_q-h_q}^q) = 0$$

whenever $k_1 + \dots + k_s + h_1 + \cdots + h_q \leq i$, $-n_j \leq k_j \leq 0$ for any $j = 1, \dots, s$ and $-m_l \leq -h_l \leq 0$ for any $l = 1, \dots, q$ except when $k_1 = \dots = k_s = 0$ and $h_l = m_l - 1$ for any $l = 1, \dots, q$.

2. E is a direct sum of bundles $\mathcal{O}$ and $\mathcal{O}(0, \dots, 0) \boxtimes \Sigma_* \boxtimes \dots \boxtimes \Sigma_*$ with some balanced twist $(t, \dots, t)$.

Proof. (1) $\Rightarrow$ (2). Let assume that t is an integer such that $E(t, \dots, t)$ is regular but $E(t-1, \dots, t-1)$ not.

By the definition of regularity and (1) we can say that $E(t-1, \dots, t-1)$ is not regular if and only if one of the following conditions is satisfied:

- i $H^d(E(t-1, \dots, t-1) \otimes \mathcal{O}(-n_1, \dots, -n_s, -m_1+1, \dots, -m_q+1)) \neq 0$.
- ii $H^{n_1+\dots+n_s+m_1-1+\dots+m_q-1}(E(t-1, \dots, t-1) \otimes \mathcal{O}(-n_1, \dots, -n_s) \boxtimes \Sigma_*(-m_1+1) \boxtimes \dots \boxtimes \Sigma_*(-m_q+1)) \neq 0$.

Let us consider one by one the conditions:

(i) Let $H^d(E(t-1, \dots, t-1) \otimes \mathcal{O}(-n_1, \dots, -n_s, -m_1+1, \dots, -m_q+1)) \neq 0$, we can conclude that $\mathcal{O}(t, \dots, t)$ is a direct summand as in the above theorem.

(ii) Let $H^{n_1+\dots+n_s+m_1-1+\dots+m_q-1}(E(t, \dots, t) \otimes \mathcal{O}(-n_1-1, \dots, -n_s-1) \boxtimes \Sigma_*(-m_1) \boxtimes \dots \boxtimes \Sigma_*(-m_q)) \neq 0$. Let us consider the following exact sequences tensored by $E(t, \dots, t)$:

$$\begin{aligned}
0 &\rightarrow \mathcal{O}(-n_1-1, \dots, -n_s-1) \boxtimes \Sigma_*(-m_1) \boxtimes \dots \boxtimes \Sigma_*(-m_q) \rightarrow \dots \\
&\dots \rightarrow \mathcal{O}(0, -n_2-1, \dots, -n_s-1) \boxtimes \Sigma_*(-m_1) \boxtimes \dots \boxtimes \Sigma_*(-m_q) \rightarrow 0, \\
0 &\rightarrow \mathcal{O}(0, -n_2-1, \dots, -n_s-1) \boxtimes \Sigma_*(-m_1) \boxtimes \dots \boxtimes \Sigma_*(-m_q) \rightarrow \dots \\
&\dots \rightarrow \mathcal{O}(0, 0, -n_3-1, \dots, -n_s-1) \boxtimes \Sigma_*(-m_1) \boxtimes \dots \boxtimes \Sigma_*(-m_q) \rightarrow 0, \\
&\dots \\
0 &\rightarrow \mathcal{O}(0, \dots, 0, -n_s-1) \boxtimes \Sigma_*(-m_1) \boxtimes \dots \boxtimes \Sigma_*(-m_q) \rightarrow \dots \\
&\dots \rightarrow \mathcal{O}(0, \dots, 0) \boxtimes \Sigma_*(-m_1) \boxtimes \dots \boxtimes \Sigma_*(-m_q) \rightarrow 0.
\end{aligned}$$

Since all the bundles in the above sequences are

$$\mathcal{E}_{n_1-k_1}^1 \boxtimes \dots \boxtimes \mathcal{E}_{n_s-k_s}^s \boxtimes \mathcal{G}_{m_1-h_1}^1 \boxtimes \dots \boxtimes \mathcal{G}_{m_q-h_q}^q$$

with decreasing indexes, by using the vanishing conditions in (1) we can see that there is a surjection from

$$H^{m_1-1+\dots+m_q-1}(E(t, \dots, t) \otimes \mathcal{O}(0, \dots, 0) \boxtimes \Sigma_*(-m_1) \boxtimes \dots \boxtimes \Sigma_*(-m_q))$$

to

$$H^{n_1+\dots+n_s+m_1-1+\dots+m_q-1}(E(t, \dots, t) \otimes \mathcal{O}(-n_1-1, \dots, -n_s-1) \boxtimes \Sigma_*(-m_1) \boxtimes \dots \boxtimes \Sigma_*(-m_q)).$$

Let us consider now the following exact sequences on $\mathcal{Q}_{m_1} \times \dots \times \mathcal{Q}_{m_q}$ for any integer p :

$$0 \rightarrow \Sigma_*(-m_1) \boxtimes \dots \boxtimes \Sigma_*(p-1) \rightarrow \Sigma_*(-m_1) \boxtimes \dots \boxtimes \Sigma_*(p)^{2(\lfloor m_q+1/2 \rfloor)} \rightarrow \Sigma_*(-m_1) \boxtimes \dots \boxtimes \Sigma_*(p) \rightarrow 0.$$

We get the long exact sequence

$$\begin{aligned}
0 &\rightarrow \Sigma_*(-m_1) \boxtimes \dots \boxtimes \Sigma_*(-m_q) \rightarrow \Sigma_*(-m_1) \boxtimes \dots \boxtimes \mathcal{O}(-m_q+1)^{2(\lfloor m_q+1/2 \rfloor)} \rightarrow \dots \\
&\dots \rightarrow \Sigma_*(-m_1) \boxtimes \dots \boxtimes \Sigma_*(-1) \rightarrow 0.
\end{aligned}$$

In the same way we can get

$$\begin{aligned}
0 \rightarrow \Sigma_*(-m_1) \boxtimes \cdots \boxtimes \Sigma_*(-1) &\rightarrow \Sigma_*(-m_1) \boxtimes \cdots \boxtimes \mathcal{O}(-m_{q-1}+1)^{2([m_{q-1}+1/2])} \rightarrow \cdots \\
&\cdots \rightarrow \boxtimes \Sigma_*(-m_1) \boxtimes \cdots \boxtimes \Sigma_*(-1) \boxtimes \Sigma_*(-1) \rightarrow 0, \\
&\cdots \\
0 \rightarrow \Sigma_*(-m_1) \boxtimes \Sigma_*(-1) \boxtimes \cdots \boxtimes \Sigma_*(-1) &\rightarrow \mathcal{O}(-m_1+1)^{2([m_1+1/2])} \boxtimes \Sigma_*(-1) \boxtimes \boxtimes \Sigma_*(-1) \rightarrow \cdots \\
&\cdots \rightarrow \Sigma_*(-1) \boxtimes \cdots \boxtimes \Sigma_*(-1) \rightarrow 0.
\end{aligned}$$

Then on $\mathbf{P}^{n_1} \times \cdots \times \mathbf{P}^{n_s} \times \mathcal{Q}_{m_1} \times \cdots \times \mathcal{Q}_{m_q}$ we can obtain the following exact sequence tensored by $E(t, \dots, t)$:

$$\begin{aligned}
0 \rightarrow \mathcal{O}(0, \dots, 0) \boxtimes \Sigma_*(-m_1) \boxtimes \cdots \boxtimes \Sigma_*(-m_q) &\rightarrow \cdots \\
&\cdots \rightarrow \mathcal{O}(0, \dots, 0) \boxtimes \Sigma_*(-1) \boxtimes \cdots \boxtimes \Sigma_*(-1) \rightarrow 0.
\end{aligned}$$

By using the vanishing conditions in (1) as above we can see that there is a surjection from

$$H^0(E(t, \dots, t) \otimes \mathcal{O}(0, \dots, 0) \boxtimes \Sigma_*(-1) \boxtimes \cdots \boxtimes \Sigma_*(-1))$$

to

$$H^{m_1-1+\cdots+m_q-1}(E(t, \dots, t) \otimes \mathcal{O}(0, \dots, 0) \boxtimes \Sigma_*(-m_1) \boxtimes \cdots \boxtimes \Sigma_*(-m_q))$$

and we can conclude that

$$H^0(E(t, \dots, t) \otimes \mathcal{O}(0, \dots, 0) \boxtimes \Sigma_*(-1) \boxtimes \cdots \boxtimes \Sigma_*(-1)) \neq 0.$$

This means that there exists a non zero map

$$f : E(t, \dots, t) \rightarrow \mathcal{O}(0, \dots, 0) \boxtimes \Sigma_* \boxtimes \cdots \boxtimes \Sigma_*.$$

On the other hand

$$\begin{aligned}
H^{n_1+\cdots+n_s+m_1-1+\cdots+m_q-1}(E(t, \dots, t) \otimes \mathcal{O}(-n_1-1, \dots, -n_s-1) \boxtimes \Sigma_*(-m_1) \boxtimes \cdots \boxtimes \Sigma_*(-m_q)) &\cong \\
&\cong H^q(E(t, \dots, t) \otimes \mathcal{O}(0, \dots, 0) \boxtimes \Sigma_*(-1) \boxtimes \cdots \boxtimes \Sigma_*(-1)).
\end{aligned}$$

Let us consider the following exact sequences tensored by $E^\vee(-t, \dots, -t)$:

$$\begin{aligned}
0 \rightarrow \mathcal{O}(0, \dots, 0) \boxtimes \Sigma_*(-1) \boxtimes \cdots \boxtimes \Sigma_*(-1) &\rightarrow \cdots \\
&\cdots \rightarrow \mathcal{O}(0, \dots, 0) \boxtimes \Sigma_* \boxtimes \cdots \boxtimes \Sigma_* \rightarrow 0.
\end{aligned}$$

By using the Serre duality and the vanishing conditions in (1) we can conclude that

$$H^0(E^\vee(-t, \dots, -t) \otimes \mathcal{O}(0, \dots, 0) \boxtimes \Sigma_* \boxtimes \cdots \boxtimes \Sigma_*) \neq 0.$$

This means that there exists a non zero map

$$g : \mathcal{O}(0, \dots, 0) \boxtimes \Sigma_* \boxtimes \cdots \boxtimes \Sigma_* \rightarrow E(t, \dots, t).$$

Then, by arguing as in [3] Theorem 1.2, we see that the composition of the maps f and g is not zero so must be the identity and we have that $\mathcal{O}(0, \dots, 0) \boxtimes \Sigma_* \boxtimes \cdots \boxtimes \Sigma_*$ is a direct summand of $E(t, \dots, t)$.

By iterating these arguments we get (2).

(2) $\Rightarrow$ (1). We argue as in Theorem 4.11. Since $H^i(\mathcal{Q}_n, \Sigma_*(e)) \neq 0$ if and only if $i = 0$ and $e \geq 0$ or $i = n$ and $e \leq -n - 1$, we have that $\mathcal{O}(0, \dots, 0) \boxtimes \Sigma_* \boxtimes \cdots \boxtimes \Sigma_*$ satisfies all the conditions in (1). $\square$

References

- [1] V. ANCONA AND G. OTTAVIANI, *Some applications of Beilinson's theorem to projective spaces and quadrics*, 1991, Forum Math. 3, no. 2, 157-176.
- [2] E. BALLICO AND F. MALASPINA, *Qregularity and an Extension of Evans-Griffiths Criterion to Vector Bundles on Quadrics*, 2007, preprint arXiv:0802.0451v1.
- [3] E. BALLICO AND F. MALASPINA, *Regularity and Cohomological Splitting Conditions for Vector Bundles on Multiprojective Spaces*, 2008, preprint arXiv:0802.0960.
- [4] A. A. BEILINSON, *Coherent sheaves on $\mathbb{P}^n$ and Problems of Linear Algebra*, 1979, Funkt. Anal. Appl. 12, 214-216.
- [5] L. COSTA AND R.M. MIRÓ-ROIG, *Tilting sheaves on toric varieties*, 2004, Math. Z. 248, 849-865.
- [6] L. COSTA AND R.M. MIRÓ-ROIG, *Cohomological characterization of vector bundles on multiprojective spaces*, 2005, J. of Algebra, 294, 73-96.
- [7] L. COSTA AND R.M. MIRÓ-ROIG, *Monads and regularity of vector bundles on projective varieties*, 2007, Preprint.
- [8] J. W. HOFFMAN AND H. H. WANG, *Castelnuovo-Mumford regularity in biprojective spaces*, Adv. Geom. 4 (2004), no. 4, 513-536.
- [9] G. HORROCKS, *Vector bundles on the punctured spectrum of a ring*, 1964, Proc. London Math. Soc. (3) 14, 689-713.
- [10] M. M. KAPRANOV, *On the derived category of coherent sheaves on Grassmann manifolds*, 1985, Math. USSR Izvestiya, 24, 183-192.
- [11] M. M. KAPRANOV, *On the derived category of coherent sheaves on some homogeneous spaces*, 1988, Invent. Math., 92, 479-508.
- [12] H. KNÖRRER, *Cohen-Macaulay modules of hypersurface singularities I*, 1987, Invent. Math. 88, 153-164.
- [13] F. MALASPINA, *A Few Splitting Criteria for Vector Bundles*, 2007, arXiv:0802.1062, Ricerche di Matematica to appear.
- [14] D. MUMFORD, *Lectures on curves on an algebraic surface*, Princeton University Press, Princeton, N.J., 1966.
- [15] G. OTTAVIANI, *Criteres de scindage pour les fibres vectoriel sur les grassmanniennes et les quadriques*, 1987, C. R. Acad. Sci. Paris, 305, 257-260.
- [16] G. OTTAVIANI, *Spinor bundles on Quadrics*, 1988, Trans. Am. Math. Soc., 307, no 1, 301-316.
- [17] G. OTTAVIANI, *Some extension of Horrocks criterion to vector bundles on Grassmannians and quadrics*, 1989, Annali Mat. Pura Appl. (IV) 155, 317-341.